



VI  
<sup>2</sup> P<0.05  
 2  
 2.1  
 73 91.3%  
 13 15.7% 10 12.0%  
 15 18.1% 4 cm 9  
 10.8% 7 8.4%  
 10 12.0%  
 1 1.2% 40 48.2% VI

2.2 VI  
 83 VI  
 VI  
 P>0.05 1

1				%	<sup>2</sup>	P
10	3	76.9	5.1	<0.05		
30	40	42.9				
45	20	44.4	5.94	<0.05		
45	27	71.1				
24	36	40	5.85	<0.05		
10	3	76.9				
34	39	46.6	0.21	>0.05		
6	4	60				
13	2	86.7	10.85	<0.01		
27	41	39.7				
4	32	43.2	4.99	<0.05		
4	8	88.9				

2.3  
 23 27.7%  
 4.8% 1 1.2%  
 D<sub>3</sub>  
 1

2.4  
 3 3.6%  
 II III IV V  
 1 1.2% PTC

3  
 3.1 PTC  
 20  
 2012  
 PTC

PTC  
 2009  
 American Thyroid Association ATA

<1 cm PTC  
 [2] 2012  
 4 cm  
 + [6]  
 42 +  
 1~3  
 PTC

3.2 CLND  
 CLND  
 CLND  
 cN<sub>0</sub> PTC  
 PTC

[7]  
 CLND  
 [8] VI

VI  
 [9-10], cN<sub>0</sub> PTC  
 , 50%~60%,  
 [11-12]

VI  
 [13-15]

VI  
 CLND  
 cN<sub>0</sub>

cN<sub>0</sub> PTC  
 PTC  
 48.2%

VI  
 VI  
 VI

CLND

cN<sub>0</sub>

### 3.3 CLND

CLND

TNM

[1] Jemal A, Siegel R, Xu J, et al. Cancer statistics, 2010[J]. CA Cancer J Clin, 2010, 60(5):277-300

[2] American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer1, Cooper DS, Doherty GM, et al. A Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer[J]. Thyroid, 2009, 19(11):1167-1264

[3] Rajeev P, Ahmed S, Ezzat TM, et al. The number of positive lymph nodes in the central compartment has prognostic impact in papillary thyroid cancer[J]. Langenbecks Arch Surg, 2013, 398(3):377-382

[4] Leboulleux S, Girard E, Rose M, et al. Ultrasound criteria of malignancy for cervical lymph nodes in patients followed up for differentiated thyroid cancer[J]. J Clin Endocrinol Metab, 2007, 92(9):3590-3594

[5] , , . cN<sub>0</sub> [J]. , 2014, 29(3):195-198

[6] . [J]. , 2012, 39(17):1249-1272

[7] Kefeli M, Akpolat I, Yildirim A, et al. Morphometric analysis in cytologic evaluation of papillary thyroid carcinoma[J]. Anal Quant Cytol Histol, 2010, 32(4): 234-238

[8] , , . Meta [J]. , 2013, 22(5):608-612

[9] Zhang L, Wei WJ, Ji QH, et al. Risk factors for neck nodal metastasis in papillary thyroid microcarcinoma: a study of 1066 patients[J]. J Clin Endocrinol Metab, 2012, 97(4):1250-1257

[10] , . ( 572 ) [J]. , 2008, 28(5):365-367.

[11] , , , . cN<sub>0</sub>

[16]

[17]

